## COLON AND RECTUM STAGING FORM

CLINICAL Extent of disease before any treatment	STAGE CATEGO	PATHOLOGIC Extent of disease through completion of definitive surgery		
y clinical – staging completed after neoadjuvant therapy but before subsequent surgery	Tumor Size:	LATERALITY: ☐ left ☐ right ☐ bilateral	y pathologic – staging completed after neoadjuvant therapy AND subsequent surgery	
□ TX □ T0 □ Tis □ T1 □ T2 □ T3 □ T4a □ T4b	Primary tumor cannot be assessed No evidence of primary tumor Carcinoma in situ: intraepithelial or invasion Tumor invades submucosa Tumor invades muscularis propria Tumor invades through the muscularis pro Tumor penetrates to the surface of the vis Tumor directly invades or is adherent to o  *Note: Tis includes cancer cells confined with (intraepithelial) or mucosal lamina prop through the muscularis mucosae into the  **Note: Direct invasion in T4 includes invasion colorectum as a result of direct extensi microscopic examination (for example, carcinoma of the cecum) or, for cancer location, direct invasion of other organs beyond the muscularis propria (i.e., res the descending colon invading the left distal rectal cancer with invasion of pro  ***Note: Tumor that is adherent to other organs However, if no tumor is present in the a should be pT1-4a depending on the an L classifications should be used to ider lymphatic invasion whereas the PN site perineural invasion.	□ TX □ T0 □ Tis □ T1 □ T2 □ T3 □ T4a □ T4b		
NX	REGIONAL LYMP Regional lymph nodes cannot be assesse No regional lymph node metastasis Metastasis in 1 to 3 regional lymph nodes Metastasis in 1 regional lymph nodes Metastasis in 2-3 regional lymph nodes Tumor deposit(s) in the subserosa, mese pericolic or perirectal tissues without of Metastasis in 4 or more regional lymph nodes Metastasis in 4 to 6 regional lymph nodes Metastasis in 7 or more regional lymph nodes Metastasis in 7 or more regional lymph nodes Metastasis in 7 or more regional lymph nodes Mote: A satellite peritumoral nodule in the peric carcinoma without histologic evidence of resid represent discontinuous spread, venous invas totally replaced lymph node (N1/2). Replaced of positive nodes in the N category, whereas disc should be classified and counted in the Site-Si (TD).	□ NX □ N0 □ N1 □ N1a □ N1b □ N1c □ N2 □ N2a □ N2b		
HOSPITAL NAME/ADDRESS		PATIENT NAME/INFORMATION		

(continued on next page)

COLON AND RECTUM STAGING FORM															
DISTANT METASTASIS (M)  No distant metastasis (no pathologic M0; use clinical M to complete stage group) Distant metastasis M1a Metastasis confined to one organ or site (e.g., liver, lung, ovary, non-regional node). M1b Metastases in more than one organ/site or the peritoneum.								0	M1 M1a M1b						
					ANATON	nic S	TAGE • P	ROGI	IOSTI	c Group	PS				
		_		CLINICA						_		OLOGIC			
	<b>OUP</b> 0	<b>T</b> Tis	<b>N</b> N0	<b>M</b> M0	Dukes*	MAC	,*	GR	<b>OUP</b> 0	<b>T</b> Tis	<b>N</b> N0	M M0	Dukes*	MAC*	
ä	I	T15	NO NO	M0	A	A			I	T1	NO NO	M0	A	A	
_		T2	N0	M0	A	B1				T2	N0	M0	Α	B1	
	IIA IIB	T3 T4a	N0 N0	M0 M0	B B	B2 B2			IIA IIB	T3 T4a	N0 N0	M0 M0	B B	B2 B2	
	IIC	T4b	NO NO	M0	В	B3			IIC	T4b	NO NO	M0	В	B3	
ā	IIIA	T1-T2	N1/N1c	M0	Č	C.			IIIA	T1-T2	N1/N1c	M0	Č	C1	
		T1	N2a	M0	С	C.				T1	N2a	M0	С	C1	
	IIIB	T3-T4a	N1/N1c	M0	С	C			IIIB	T3-T4a	N1/N1c	M0	С	C2	
		T2-T3 T1-T2	N2a N2b	M0 M0	C	C.	1/C2 1			T2-T3 T1-T2	N2a N2b	M0 M0	C C	C1/C2 C1	
	IIIC	T4a	N2a	M0	Č	C			IIIC	T4a	N2a	M0	Č	C2	
		T3-T4a	N2b	M0	С	C2				T3-T4a	N2b	M0	С	C2	
	11.7.4	T4b	N1-N2	M0	С	C	3		11.7.4	T4b	N1-N2	M0	С	C3	
	IVA IVB	Any T Any T	Any N Any N	M1a M1b	-	-			IVA IVB	Any T Any T	Any N Any N	M1a M1b	-	-	
*Dukes B is a composite of better (T3 N0 M0) and worse (T4 N0 M0) prognostic  *Dukes B is a composite of better (T3 N0 M0) and worse (T4 N0 M0) prognostic										and worse (T	4 N0 M0)				
		Dukes C (Al classification		and Any	T N2 M0). M	AC is th	ne modified						and Any T	N2 M0). MAC	is
	Stage ur		1.						the modified Astler-Coller classification.  ☐ Stage unknown						
	Jiago ai	iitiiowii							Olago o						
			PROGN	IOSTIC F	ACTORS (	SITE-	SPECIFIC FA	CTOR	S)			General	Notes:		
REC	UIRED	FOR STA	GING: No	ne										pecial cases of	
		Y SIGNIFIC					(0-1)					TNM or pTNM classifications, the "m" suffix and "y," "r," and "a" prefixes are used. Although they do not affect the stage grouping, they indicate cases			
		ative or pro eposits (TI		nt carcino	-	-	n (CEA) ng/m				-				
	Circumfe	rential Res	section Ma	rain (CR	M)						'				
Circumferential Resection Margin (CRM) needing separate analysis.									-						
Microsatellite Instability (MSI)								m suffix indicates the presence of multiple primary tumors in a single							
Tumor Regression Grade (with neoadjuvant therapy) KRAS gene analysis									site and is recorded in parentheses:						
18g loss of heterozygosity (LOH) assay									pT(m)NN	1.					
y prefix indicates those cases in															
Hist	Histologic Grade (G) (also known as overall grade)  Grading system Grade											tial multimodali	lity		
		<i>ng systen</i> le system	1				Grade I or 1						The cTNM or		•
	-	de system					Grade II or 2							by a "y" prefix. I categorizes	
	•	-					Grade III or 3					the exten	t of tumor ac	tually present	
	-	de system	do ouetem	, io ovoilo	blo									nation. The "y' an estimate of	
	NO ∠,	3, or 4 gra	de system	ı is avalla	ible	<u> </u>	Grade IV or 4							dality therapy.	
HOSPITAL NAME/ADDRESS					Рат	PATIENT NAME/INFORMATION									

(continued from previous page)

## COLON AND RECTUM STAGING FORM


		General Notes (continued):		
ADDITIONAL DESCRIPTORS  Lymphatic Vessel Invasion (L) and Venous Invasion (V) have been convasion (LVI) for collection by cancer registrars. The College of American should be used as the primary source. Other sources may be used in the	r prefix indicates a recurrent tumor when staged after a disease-free interval, and is identified by the "r" prefix: rTNM.			
is given to positive results.	a prefix designates the stage determined at autopsy: aTNM.			
<ul> <li>□ Lymph-Vascular Invasion Not Present (absent)/Not Identified</li> <li>□ Lymph-Vascular Invasion Present/Identified</li> <li>□ Not Applicable</li> <li>□ Unknown/Indeterminate</li> </ul> Residual Tumor (R)	surgical margins is data field recorded by registrars describing the surgical margins of the resected primary site specimen as determined only by the pathology report.			
The absence or presence of residual tumor after treatment. In some case with neoadjuvant therapy there will be residual tumor at the primary site a incomplete resection or local and regional disease that extends beyond the	neoadjuvant treatment is radiation therapy or systemic therapy (consisting of chemotherapy,			
<ul> <li>□ RX Presence of residual tumor cannot be assessed</li> <li>□ R0 No residual tumor</li> <li>□ R1 Microscopic residual tumor</li> <li>□ R2 Macroscopic residual tumor</li> </ul>	hormone therapy, or immunotherapy) administered prior to a definitive surgical procedure. If the surgical procedure is not performed, the administered therapy no longer meets the definition of neoadjuvant therapy.			
☐ Clinical stage was used in treatment planning (describe):				
□ National guidelines were used in treatment planning □ NCCN	Other (describe):			
Physician signature		nte/Time		
HOSPITAL NAME/ADDRESS	PATIENT NAME/INFORMATION			

(continued on next page)

## COLON AND RECTUM STAGING FORM

## Illustration

Indicate on diagram primary tumor and regional nodes involved.



HOSPITAL NAME/ADDRESS

PATIENT NAME/INFORMATION

(continued from previous page)