Nuclear Medicine MIBG Study:



Name:	Injection Date:	Check In Time:	
	Scan #1 Date:	Check In Time:	
	Scan #2 Date:	Check In Time:	
	Scan #3 Date:	Check In Time:	

	2 Days Before MIBG Injection	1 Day Before MIBG Injection	Day of MIBG Injection	Days of Each Scan
Ŗ	Your physician will prescribe the medication, potassium lodide Solution. This will	R Take first dose of Potassium lodide solution as directed	R Take second dose of Potassium lodide solution as directed	You will take pictures on three separate days
	protect your thyroid gland. Fill your prescription at the local pharmacy.	Take all medications as prescribed by your physician	Take all medications as prescribed by your physician	Wear loose comfortable clothing free from metal objects
	Take all medications as prescribed by your physician	Hydrate - drink cold beverages throughout the day	Hydrate - drink cold beverages throughout the day	R Continue to take each morning for another five (5) days, the Potassium lodide Solution
2	If you are breast-feeding, please call Nuclear Medicine at (206) 223-6951 for additional instructions	Eat meals unless you have another test that has dietary restrictions	Eat meals unless you have another test that has dietary restrictions	Hydrate - drink cold beverages throughout the day
			Female patients, who might be pregnant, e.g. last menstrual cycle greater than 28 days, will require a pregnancy test	Eat meals unless you have another test that has dietary restrictions