


# Nuclear Medicine Renal Captopril Study:












Name: \_\_\_\_\_

Appointment date: \_\_\_\_\_

Check In Time: \_\_\_\_\_

If you take medications for diabetes, call your physician or Nuclear Medicine at (206) 223-6951 for special fasting guidelines. Otherwise, please follow instructions below:

2 Days Before Appointment		Day of Appointment
 <p>Discontinue Angiotensin II type medications until after the exam. These include:</p> <ul style="list-style-type: none"> <li>Aceon, Accupril</li> <li>Altace, Atacand</li> <li>Capoten, Capozide</li> <li>Cozaar, Diovan</li> <li>Lexxel, Lotensil</li> <li>Lotensin, Lotrel</li> <li>Mavik, Micardis</li> <li>Monopril, Prinivil</li> <li>Tarka, Teveten</li> <li>Uniretic, Univasc</li> <li>Vaseretic, Vasotec</li> <li>Zestoretic, Zestoretic</li> <li>Zestril</li> </ul>	<p> Take all other medications prescribed by your physician</p> <p> If you are breast-feeding or taking diabetic medication, call Nuclear Medicine at (206) 223-6951 for special instructions</p> <p> <b>Eat and drink</b> unless you have another test that has dietary restrictions</p> <p> <b>Hydrate</b> - drink cold beverages throughout the day</p>	<p> <b>DO NOT</b> eat solid food 4 hours before your appointment</p> <p> <b>Drink</b> 2 cups of water during the hours before your appointment</p> <p><b>Drink</b> 1 cup of water when you arrive in Nuclear Medicine</p> <p> Wear loose comfortable clothing free from metal objects</p> <p> Female patients, who might be pregnant, e.g. last menstrual cycle greater than 28 days, will require a pregnancy test</p>